

## Problem set - 4

Due: Nov. 11th by 5pm in TA box.

- Use the Boltzmann equation to find the thermoelectric coefficient  $\lambda$  for a dilute (free) electron gas - which describe the situation in a semiconductor. Assume that  $\mu$  is constant while while temperature has a finite gradient for your calculation. For your final answer, recall that for a dilute gas the chemical potential is  $\mu = T \ln(n\lambda_T^3)$ , where  $\lambda_T = \frac{h}{\sqrt{2\pi m T}}$ , and  $n\lambda_T^3 \ll 1$ . Note that this applies to each of the spin components separately. Assume that the density  $n$  as well as temperature  $T$ , mass  $m$ , and scattering time  $\tau$  are given. Hint: The distribution of a dilute electron gas is still the Fermi-Dirac distribution, but with the exponential term dominating the denominator, and thus resulting in a Boltzmann distribution. So one can use the derivaton in class for  $\delta f$ , but then the current calculation must be reconsidered, as there is no onger a  $\delta$ -function in the integral.

- Phonons.

Consider a chain made out of a unit cell with two atoms in each unit cell. The atoms can move only in one direction, and have masses  $m_A$  and  $m_B$ . If the distance between the atoms change, there is an elastic energy cost of  $E_{pot} = \frac{1}{2}k(\Delta x - a/2)^2$  with  $a$  being the lattice constant, and  $a/2$  the distance between the two atoms.

- What is the dispersion of phonons in the chain? Identify an optical and acoustic branch.
- What is the speed of sound?
- What is the heat capacity of the system at low temperatures?

Note that in the limit of  $m_A = m_B$  we obtain a chain with lattice constant  $a/2$ .

- Tight binding approximation preview.

Consider a chain of  $N$  atoms, with electrons described by an Hamiltonian  $H$ . We would like to find the lowest lying states of this hamiltonian using the ground state orbital in each atom. Mark the lowest orbital in the atom  $i$  as  $|i\rangle$ .

By minimizing the energy of the states, show that the best lowest-energy  $N$  solutions that are superpositions of the form  $|\psi\rangle = \sum_{i=1}^N \alpha_i |i\rangle$  have energies that given by eigenvalues of the matrix:

$$H_{eff} = M^{-1/2} h M^{-1/2} \tag{1}$$

with  $M_{ij} = \langle i|j\rangle$ , and  $h = \langle i|H|j\rangle$ . How are the corresponding eigenstates relate to the  $|\psi\rangle$  states and the  $\alpha$ 's defined above?

In a tight-binding approximation we assume that  $h_{ij}$  and  $M_{ij}$  are nonzero only if  $|i - j| \leq 1$ .